

Subject Code: R13207/R13

Set No - 1

I B. Tech II Semester Regular Examinations August - 2014

**MATHEMATICS-II (MATHEMATICAL METHODS)**

(Common to CE, ME, CSE, PCE, IT, Chem E, Aero E, Auto E, Min E, Pet E, Metal E)

Time: 3 hours

Max. Marks: 70

Question Paper Consists of **Part-A** and **Part-B**  
Answering the question in **Part-A** is Compulsory,  
Three Questions should be answered from **Part-B**

\*\*\*\*\*

**PART-A**

- 1.(i) Write iterative scheme to find the  $n^{\text{th}}$  root of a real number  $K(>0)$ .
- (ii) Find  $\Delta \log f(x)$ .
- (iii) Find half range Fourier sine series of  $f(x) = e^x$  in  $(0, 1)$ .
- (iv) Prove that  $Z(\sinh nt) = \frac{z \sinh t}{z^2 - 2z \cosh t + 1}$ .
- (v) Using Euler's method, find the value of  $y(0.5)$  (take  $h = 0.25$ ) and compare with the exact solution of the equation  $y' = x + y$ ,  $y(0) = 1$
- (vi) If  $F_p$  is complex Fourier transform of  $f(x)$ , then find the complex Fourier transform of  $f(x) \sin ax$ .

[3+3+3+3+5+5]

**PART - B**

- 2.(a) Using Newton-Raphson method find the root of the equation  $x + \log_{10} x = 3.375$  correct to four decimal places.
- (b) The population of a town in the decimal census is given below. Estimate the population of a town for the year 1895

Year X	1971	1981	1991	2001	2011
Population Y	146	166	181	193	201

[8+8]

- 3.(a) Find positive root of  $x^3 - 5x + 3 = 0$  using Regula falsi method up to 4 steps.
- (b) Using Lagrange's interpolation formulae find the value of  $y(12)$  from the data

X	5	7	9	13
Y	11	13	18	27

[8+8]

- 4.(a) Solve  $y' = x^2 y + 1$ ,  $y(0)=1$  using Taylors method up to 3<sup>rd</sup> degree term and compute  $y(0.1)$ .
- (b) Find the fourier series of  $f(x) = x \sin x$  in  $(-\pi, \pi)$ .

[8+8]



5.(a) Find half range cosine series of  $f(x) = \begin{cases} 1, & 0 < x < \frac{\pi}{2} \\ -1, & \frac{\pi}{2} < x < \pi \end{cases}$ .

(b) Use Runge-Kutta 4<sup>th</sup> to compute  $y(1.25)$  given that  $\frac{dy}{dx} = \frac{x^2 + y}{x}$ ,  $y(1) = 2$

[8+8]

6.(a) Find Fourier transform  $f(x) = \begin{cases} x & \text{if } |x| \leq 1 \\ 0 & \text{if } |x| > 1 \end{cases}$ .

(b) Find Z-transform of  $n a^n$ .

[8+8]

7.(a) Find Fourier sine transform of  $e^{-x}$  and hence deduce the inversion formula.

(b) Solve the difference equation  $u_{n+2} - u_n = 2^n$ ,  $u_0 = 0$ ,  $u_1 = 1$ , using Z- transforms.

[8+8]



**I B. Tech II Semester Regular Examinations August - 2014**  
**MATHEMATICS-II (MATHEMATICAL METHODS)**  
 (Common to CE, ME, CSE, PCE, IT, Chem E, Aero E, Auto E, Min E, Pet E, Metal E)

**Time: 3 hours**

**Max. Marks: 70**

Question Paper Consists of **Part-A** and **Part-B**  
 Answering the question in **Part-A** is Compulsory,  
 Three Questions should be answered from **Part-B**

\*\*\*\*\*

**PART-A**

- 1.(i) Using bisection method find the first four approximations to the real root of  $3x = e^x$ .
- (ii) Prove that  $\Delta\left(\frac{1}{f(x)}\right) = \frac{-\Delta f(x)}{f(x)f(x+1)}$ .
- (iii) If  $Z(n^2) = \frac{z^2+z}{(z-1)^3}$  find  $Z(n^3)$ .
- (iv) Find the Half range Fourier sine series of  $f(x) = |x|$  in  $(0, 1)$ .
- (v) If  $y' = 2x - y$ ,  $y(1) = 3$ , find the solution, up to third degree term, using Picard's method.
- (vi) Prove  $F[x^n f(x)] = (-i)^n \frac{d^n}{dp^n} [F(p)]$ .

[3+3+3+3+5+5]

**PART - B**

- 2.(a) Using Newton – Raphson method, find a root of the equation  $2x - 3\sin x = 5$  near  $x=5$  correct to three decimal places.
  - (b) Given that  $f(6500) = 80.8084$ ,  $f(6510) = 80.6846$ ,  $f(6520) = 80.7456$ ,  $f(6530) = 80.8084$ , find  $f(6526)$  using Gauss backward interpolation formula. [8+8]
  - 3.(a) Find a positive root of  $2x = 3 + \cos x$  by using Newton-Raphson method correct to three decimal places. (Use Bisection method for the first approximation).
  - (b) Using Lagrange's Interpolation formula for the value of  $y(6)$  given the following table
- |   |      |      |      |     |
|---|------|------|------|-----|
| X | 1    | 2.5  | 5    | 7   |
| Y | 2.25 | 4.13 | 7.25 | 9.0 |
- [8+8]
- 4.(a) Solve  $y' = y + x$ ,  $y(0) = 1$  using Picard's method up to third approximation and hence find the value of  $y(0.1)$ .
  - (b) Find the Fourier expansion of  $f(x) = x \cos x$ ,  $0 < x < 2\pi$ . [8+8]
  - 5.(a) Find half range cosine series of  $f(x) = \begin{cases} 1, & 0 < x < 1 \\ -1, & 1 < x < 2 \end{cases}$ .
  - (b) Find  $y(0.1)$  using 4<sup>th</sup> order Runge-Kutta method given that  $y' = x + x^2 y$ ,  $y(0) = 1$ . [8+8]



6.(a) Find the Fourier transform of  $\frac{1}{\sqrt{|x|}}$ .

(b) Find Z-transform of  $n^2 e^{n\theta}$ .

[8+8]

7.(a) Find Fourier cosine transform of  $\frac{1}{1+x^2}$  and hence find Fourier sine transform of  $\frac{x}{1+x^2}$ .

(b) Solve  $y(n+2) + 3y(n+1) + 2y(n) = 0$ ,  $y(0) = 0$ ,  $y(1) = 1$  using Z-Transform.

[8+8]



**I B. Tech II Semester Regular Examinations August - 2014**  
**MATHEMATICS-II (MATHEMATICAL METHODS)**  
 (Common to CE, ME, CSE, PCE, IT, Chem E, Aero E, Auto E, Min E, Pet E, Metal E)

**Time: 3 hours**

**Max. Marks: 70**

Question Paper Consists of **Part-A** and **Part-B**  
 Answering the question in **Part-A** is Compulsory,  
 Three Questions should be answered from **Part-B**

\*\*\*\*\*

**PART-A**

- 1.(i) Find reciprocal of a real number 19 using Regula falsi method.
- (ii) Expand the shift operator E in terms of exponential function.
- (iii) Employ Taylor's method to obtain the values of y(1.1) for the differential equation  $y' = xy^{1/3}, y(1) = 1$ .
- (iv) A sinusoidal voltage  $E \cos \omega t$  is passed through a half wave rectifier which clips the negative portion of the wave. Develop the resulting periodic function

$$u(t) = \begin{cases} 0 & , -\frac{T}{2} < t < 0 \\ E \cos \omega t, & 0 < t < \frac{T}{2} \end{cases}, T = \frac{2\pi}{\omega} \text{ as Fourier series.}$$

(v) Prove that  $F_s \left[ \frac{d}{dx} F(x) \right] = -p F_c(p)$

- (vi) Find the Z-transform of  $\sin((n+1)t)$ .

[3+3+3+5+3+5]

**PART - B**

- 2.(a) By using Regula-Falsi method for a real root of  $xe^x = 2$  up to 4 stages.
- (b) Using a forward difference formula, find y(11) from the given table


X	1	6	11	16	21	26
Y	5	10	14	18	24	32

[8+8]

- 3.(a) Using Newton-Raphson formula, find the root of  $e^x - x^3 + \cos 25x = 0$  around  $x = 4.5$  correct to 3 decimal places.
- (b) Using Lagrange's Interpolation formula, find the value y(2) given the following table of values

X	1	1.1	1.4	1.8
Y	2	4	8	11

[8+8]



4.(a) Using Euler's method, solve for y (0.6) from  $y' = -2xy$ ,  $y(0) = 1$  using step size 0.2.

(b) Find the Fourier series of  $f(x) = \begin{cases} 0, & -\pi < x < 0 \\ \frac{\pi}{4}, & 0 < x < \pi \end{cases}$ .

[8+8]

5.(a) Represent the function as Fourier cosine series  $f(x) = \begin{cases} \frac{\pi}{2}, & 0 < x < \frac{\pi}{2} \\ \pi - x, & \frac{\pi}{2} < x < \pi \end{cases}$ .

(b) Use Runge-Kutta 4<sup>th</sup> order to compute y(1.2) for the equation  $y' = \frac{x^2 + y}{x}$ ,  $y(1) = 2$ .

[8+8]

6.(a) Find the Fourier cosine transform of  $\frac{e^{-ax}}{x}$ .

(b) Find  $Z^{-1} \left[ \frac{8z - z^3}{(4 - z)^3} \right]$ .

[8+8]

7.(a) Find Fourier cosine transform of  $f(x) = \begin{cases} x & \text{if } |x| \leq a \\ 0 & \text{if } |x| > a \end{cases}$ .

(b) Solve  $u_{n+2} - 6u_{n+1} + 9u_n = 0$  using Z-transform.

[8+8]



Subject Code: R13207/R13

Set No - 4

I B. Tech II Semester Regular Examinations August - 2014

**MATHEMATICS-II (MATHEMATICAL METHODS)**

(Common to CE, ME, CSE, PCE, IT, Chem E, Aero E, Auto E, Min E, Pet E, Metal E)

Time: 3 hours

Max. Marks: 70

Question Paper Consists of **Part-A** and **Part-B**  
Answering the question in **Part-A** is Compulsory,  
Three Questions should be answered from **Part-B**

\*\*\*\*\*

**PART-A**

- 1.(i) Evaluate  $\sqrt[4]{29}$  to four decimal places by Newton-Raphson method.
- (ii) If the interval of differencing is unity, find  $\Delta^2 \sin(px + q)$ .
- (iii) Using Taylor's series method obtain  $y(0.2)$  for the differential equation  $y' + 2y = 3e^{2x}$ ,  $y(0) = 0$ .
- (iv) Find the Fourier series of  $f(x) = |\cos x|$  in  $(-\pi, \pi)$ .
- (v) Find Fourier transform of  $f(x) = \begin{cases} 1, & |x| < a \\ 0, & |x| > a \end{cases}$ .
- (vi) Prove that  $Z(\cos nt) = \frac{z(z - \cos t)}{z^2 - 2z \cos t + 1}$ .

[3+3+3+3+5+5]

**PART - B**

- 2.(a) Find a real root of  $x^3 - 4x - 9 = 0$  using Bisection method up to 4 stages.
- (b) Using Gauss Backward difference polynomial, find  $y(5)$  given that

X	0	4	6	8	10
Y	5	11	13	15	17

[8+8]

- 3.(a) Using Newton-Raphson method, find a positive root of  $\cos x - x e^x = 0$  up to four decimal places.
- (b) Using Lagrange's Interpolation, find  $f(12)$ , given that

X	3	7	9	13
Y	5	12	13	21

[8+8]

- 4.(a) Using Euler's method, solve for  $y(0.4)$  from  $y' = 2xy$ ,  $y(0) = 1$  using step size 0.2.
- (b) Find the Fourier series of periodicity 2 for  $f(x) = x + x^2$  in  $0 < x < 2$ .

[8+8]



5.(a) Represent the function as Fourier sine series  $f(x) = \begin{cases} \frac{\pi}{2}, & 0 < x < \frac{\pi}{2} \\ \pi - x, & \frac{\pi}{2} < x < \pi \end{cases}$ .

(b) Estimate  $y(0.2)$ , given  $y' = 3x + y, y(0) = 1$  using Runge-Kutta 4<sup>th</sup> order.

[8+8]

6.(a) Find Fourier cosine transform of  $\frac{e^{-ax}}{x}$ .

(b) Find the Z-transform of  $\{x(n)\} = n z^n$

[8+8]

7.(a) Find Fourier transform of  $f(x) = \begin{cases} \frac{1}{2a}, & |x| \leq a \\ 0, & |x| > a \end{cases}$ .

(b) Solve  $u_{n+2} - u_n = 2^n, u_0 = 0, u_1 = 1$  using Z-transform.

[8+8]

