

Subject Code: R13207/R13

Set No - 1

I B. Tech II Semester Regular/Supply Examinations July - 2015

**MATHEMATICS-II (MATHEMATICAL METHODS)**

(Common to CE, ME, CSE, PCE, IT, Chem E, Aero E, Auto E, Min E, Pet E, Metal E)

Time: 3 hours

Max. Marks: 70

Question Paper Consists of **Part-A** and **Part-B**  
Answering the question in **Part-A** is Compulsory,  
Three Questions should be answered from **Part-B**

\*\*\*\*\*

**PART-A**

- 1.(a) Define Rate of convergence and what is the rate of convergence of bisection and iteration methods
- (b) Find  $\Delta f(x)$  if  $f(x) = \frac{2x+1}{x(x+1)}$  by taking  $h=1$
- (c) Write the merits and demerits of Taylor's Method
- (d) Find the Fourier series of  $f(x) = \sin x$  in  $(-\pi, \pi)$
- (e) Find the Fourier cosine transform of  $f(x) = 1$  in  $(0, 2)$
- (f) Find  $Z^{-1} \left[ \frac{z^2}{z^2+1} \right]$

[3+4+3+4+4+4]

**PART-B**

- 2.(a) Find the real root of  $x + \log_{10} x - 2 = 0$  using Newton Raphson method
  - (b) Find the positive root of the equation  $x^3 - 9x + 1 = 0$  by Bisection Method
- [8+8]
- 3.(a) Compute  $y^1(4)$  from following table
- |   |   |   |   |    |    |
|---|---|---|---|----|----|
| X | 1 | 2 | 4 | 8  | 10 |
| Y | 0 | 1 | 5 | 21 | 27 |
- (b) Find by Gauss's Backward interpolating formula the value of  $y$  at  $x=1936$ , using the following table:
- |     |      |      |      |      |      |      |
|-----|------|------|------|------|------|------|
| $x$ | 1901 | 1911 | 1921 | 1931 | 1941 | 1951 |
| $y$ | 12   | 15   | 20   | 27   | 39   | 52   |
- [8+8]
- 4.(a) Find  $y(2)$  and  $y(3)$  by Picard's method given that  $\frac{dy}{dx} = 2x - y$ ,  $y(1) = 3$ .
  - (b) Using Modified Euler's method of fourth order evaluate  $y(0.1)$  and  $y(0.2)$  given that  $y^1 = x + y$ ,  $y(0) = 1$ .

[8+8]



5.(a) Obtain the Fourier series for the function  $f(x)$  given by  $f(x) = \begin{cases} -x(x + \pi); -\pi \leq x \leq \pi \\ x(x + \pi); 0 \leq x \leq \pi \end{cases}$

(b) Obtain the Fourier series expansion of  $f(x)$  given that  $f(x) = (\pi - x)^2$  in  $0 < x < 2\pi$  and

Deduce the value of  $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{6}$ .

[8+8]

6.(a) Find the Fourier cosine transform of  $f(x) = \begin{cases} \cos x \text{ if } |x| < a \\ 0 \text{ if } |x| > a \end{cases}$

(b) Find the finite Fourier sine and cosine transforms of  $f(x) = e^{-ax}$  in  $(0, L)$


[8+8]

7.(a) Find  $Z^{-1}\left(\frac{2z}{z^3 - z^2 + z - 1}\right)$

(b) If  $F(z) = \frac{5z^2 + 3z + 12}{(z - 1)^4}$ ; then find the values of  $f(2)$  and  $f(3)$

[8+8]

\*\*\*\*\*



**I B. Tech II Semester Regular/Supply Examinations July - 2015**  
**MATHEMATICS-II (MATHEMATICAL METHODS)**

(Common to CE, ME, CSE, PCE, IT, Chem E, Aero E, Auto E, Min E, Pet E, Metal E)

**Time: 3 hours**

**Max. Marks: 70**

Question Paper Consists of **Part-A** and **Part-B**  
 Answering the question in **Part-A** is Compulsory,  
 Three Questions should be answered from **Part-B**

\*\*\*\*\*

**PART-A**

- 1.(a) Write the working rule to find the root of the equation by bisection method
- (b) Evaluate the expression  $(1 + \Delta)(1 - \nabla)$
- (c) Write the merits and demerits of Euler's Method
- (d) Find the Fourier series of  $f(x) = \cos ax$  in  $(-L, L)$
- (e) Find the Fourier sine transform of  $f(x) = 1$  in  $(0, \pi)$
- (f) Find  $Z^{-1} \left[ \frac{z^2}{(z-a)(z-b)} \right]$  using convolution theorem

[3+4+3+4+4+4]

**PART-B**

- 2.(a) Find the positive root of the equation  $x^3 - 5x - 7 = 0$  by False position method
- (b) Find the positive root of the equation  $e^x - 3x = 0$  by Newton Raphson method

[8+8]

- 3.(a) Find  $f(2.5)$  from the following table

x	1.6	1.8	2.0	2.2	2.4	2.6
y	4.95	6.05	7.39	9.03	11.02	13.46

- (b) Using Lagrange's formula, calculate  $f(3)$  from the table:

x	0	1	2	4	5	6
f(x)	1	14	15	5	6	19

[8+8]

- 4.(a) Using Taylor's series method: Solve  $y' = xy + y^2, y(0) = 1$  at  $x = 0.1, 0.2, 0.3$
- (b) Solve:  $y' = y - x, y(0) = 2, h = 0.2$  find  $y(0.2)$ , using R- K method.

[8+8]

- 5.(a) Develop the Fourier series of  $f(x) = \begin{cases} 2; -2 \leq x \leq 0 \\ x; 0 \leq x \leq 2 \end{cases}$

- (b) If  $f(x) = |\cos x|$ ; Expand  $f(x)$  as a Fourier series in the interval  $(-\pi, \pi)$

[8+8]



6.(a) Find Fourier Sine transform  $f(x) = \frac{1}{(x^2 + 1)}$

(b) Express the function  $f(x) = \begin{cases} 0; x < 0 \\ \frac{1}{2}; x = 0 \\ e^{-x}; x > 0 \end{cases}$  as a Fourier integral.

[8+8]

7.(a) Solve the difference equation  $y_{n+2} - 2y_{n+1} + y_n = 3n + 5$  if  $y_0 = 1, y_1 = -4$ .  
by Z -transforms

(b) Find the Z- transform of the following (i)  $n^2 e^{-an}$  (ii)  $(n + 1)^2$  (iii)  $a^n \sin(nt)$

[8+8]

\*\*\*\*\*



Subject Code: R13207/R13

Set No - 3

I B. Tech II Semester Regular/Supply Examinations July - 2015

**MATHEMATICS-II (MATHEMATICAL METHODS)**

(Common to CE, ME, CSE, PCE, IT, Chem E, Aero E, Auto E, Min E, Pet E, Metal E)

Time: 3 hours

Max. Marks: 70

Question Paper Consists of **Part-A** and **Part-B**  
Answering the question in **Part-A** is Compulsory,  
Three Questions should be answered from **Part-B**

\*\*\*\*\*

**PART-A**

- 1.(a) Write the working rule to find the root of the equation by Newton Raphson method
- (b) Find  $\left(\frac{\Delta^2}{E}\right)x^3$
- (c) Write the working rule to solve the  $y' = f(x, y)$   $y(x_0) = y_0$  by Picard's method
- (d) Find the half range sine series of  $f(x) = 2x$  in  $(0, L)$
- (e) Find the Fourier transform of  $f(x) = 1$  in  $(-1, 1)$
- (f) Find  $Z[\sin(n+1)\theta]$  using shifting theorem

[3+4+3+4+4+4]

**PART-B**

- 2.(a) Find the root of the equation:  $x^3 = 2x + 5$  by iteration method.
- (b) Find the real root for  $xe^x = 2$  by using Regula – Falsi method.

[8+8]

- 3.(a) Using Lagrange's formula fit a polynomial to the following data

x	0	1	4	5
y	4	3	24	39

- (b) Estimate  $f(1.75)$  from the following table using Newton forward interpolation formula

x	1.7	1.8	1.9	2.0
y	5.474	6.050	6.686	7.389

[8+8]

- 4.(a) Using Runge Kutta method of fourth order evaluate  $y(0.1)$  and  $y(0.2)$  given that  $y' = x + y$ ,  $y(0) = 1$

- (b) Apply Taylor series methods to find  $y(1.1)$ ,  $y(1.2)$  correct to 3 decimal places, given

$$\frac{dy}{dx} = xy^{1/3}, y(0)=1.$$

[8+8]



5.(a) Obtain the Fourier series of  $f(x) = \sqrt{1 - \cos x}$  in  $(-\pi, \pi)$

(b) Expand  $f(x) = \begin{cases} \frac{1}{4} - x; 0 \leq x \leq 1/2 \\ x - \frac{3}{4}; 1/2 \leq x \leq 1 \end{cases}$  as a Fourier series of sine terms.

[8+8]

6.(a) Find Fourier transform of  $f(x) = e^{-a|x|}$  ( $a > 0$ ) and hence show that

$$\int_0^{\infty} \frac{\cos(sx)}{a^2 + s^2} ds = \frac{\pi}{2a} e^{-a|x|}$$

(b) Find the finite Fourier cosine transform of

i)  $f(x) = \frac{x^2}{2\pi} - \frac{\pi}{6}, 0 \leq x \leq \pi$     ii)  $f(x) = x, 0 < x < 4$


[8+8]

7.(a) Using Z- Transform solve  $y_{n+1} + 2y_{n+1} + y_n = n$ ; Given that  $y_0 = y_1 = 0$ ;

(b) Using  $Z(n^2) = \frac{z^2 + z}{(z-1)^3}$  prove that  $Z((n+1)^2) = \frac{z^3 + z^2}{(z-1)^3}$

[8+8]

\*\*\*\*\*



## I B. Tech II Semester Regular/Supply Examinations July - 2015

## MATHEMATICS-II (MATHEMATICAL METHODS)

(Common to CE, ME, CSE, PCE, IT, Chem E, Aero E, Auto E, Min E, Pet E, Metal E)

Time: 3 hours

Max. Marks: 70

Question Paper Consists of **Part-A** and **Part-B**  
 Answering the question in **Part-A** is Compulsory,  
 Three Questions should be answered from **Part-B**

\*\*\*\*\*

**PART-A**

- 1.(a) What is mean by quadratic convergence and derive the convergence condition for Newton Rapson method.
- (b) Find  $\Delta f(x)$  if  $f(x) = \frac{1}{(x^2 + 5x + 6)}$  by taking  $h = 1$
- (c) Write the working rule to solve the  $y' = f(x, y)$   $y(x_0) = y_0$  by RK method of third order
- (d) Find the half range cosine series of  $f(x) = x$  in  $(0, \pi)$
- (e) Find the Finite Fourier cosine transform of  $f(x) = 1$  in  $(0, \pi)$
- (f) Find  $Z[\cos(n+1)\theta]$  using shifting theorem

[4+4+3+4+3+4]

**PART-B**

- 2.(a) Evaluate  $\sqrt{12}$  and  $\frac{1}{\sqrt{12}}$  by the fixed point iteration method.
  - (b) Find a root correct to 3 decimal places for the equation  $x^3 - 4x + 9 = 0$  using Bisection method
- [8+8]
- 3.(a) Certain values of  $x$  and  $\log_{10}^x$  are (300,2.4771),(304,2.4829),(305,2.4843),(307,2.4871).  
Find  $\log_{10}^{301}$
  - (b) Using Lagrange's Interpolation formula evaluate  $y(6)$ .

x	3	5	7	9	11
y	6	24	58	108	74

[8+8]

- 4.(a) Given  $\frac{dy}{dx} - \sqrt{xy} = 2$  and  $y(1)=1$ . Find the value of  $y(1.5)$  in steps of 0.25 using Euler's modified method.
- (b) Use Runge-Kutta method to solve  $\frac{dy}{dx} = xy + y^2, y(0) = 1$  for  $y(0.1)$  and  $y(0.2)$ .

[8+8]



5.(a) Obtain Fourier series for the function f(x) given by  $f(x) = \begin{cases} 1 + \frac{2x}{\pi}, & -\pi \leq x \leq 0 \\ 1 - \frac{2x}{\pi}, & 0 \leq x \leq \pi \end{cases}$

and deduce that  $\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots = \frac{\pi^2}{8}$

(b) If  $f(x) = \begin{cases} x; & 0 < x < \pi/2 \\ \pi - x; & \pi/2 < x < \pi \end{cases}$

Show that  $f(x) = \frac{\pi}{4} - \frac{2}{\pi} \left[ \frac{1}{1^2} \cos 2x + \frac{1}{3^2} \cos 6x + \frac{1}{5^2} \cos 10x + \dots \right]$

[8+8]

6.(a) Show that the Fourier transform of  $f(x) = \begin{cases} a - |x|, & \text{for } |x| < a \\ 0, & \text{for } |x| > a \end{cases}$  is  $\sqrt{\frac{2}{\pi}} \left( \frac{1 - \cos as}{s^2} \right)$

Hence deduce that  $\int_0^\infty \left( \frac{\sin t}{t} \right)^2 = \frac{\pi}{2}$

(b) Find the finite Fourier sine transform of f(x) defined by  $f(x) = \left( 1 - \frac{x}{\pi} \right)^2$  where  $0 < x < \pi$

[8+8]

7.(a) Solve the difference equation  $y_{n+2} - 2y_{n+1} + y_n = 2^n$  if  $y_0 = 2, y_1 = 1$ . by Z –transforms

(b) Find  $Z^{-1} \left[ \frac{z^3}{(z-3)(z^2+1)} \right]$  using the convolution theorem.

[8+8]

\*\*\*\*\*

